

Anacostia Wetlands: Has RECONSTRUCTION led to RESTORATION?

Dick Hammerschlag, Cairn Krafft, Andy Baldwin



Reconstructed Freshwater Tidal Wetlands in the Anacostia



Cahoon Guntenspergen Neckles Irwin Pattee

Marion Olsen


Twedt Beyers Dawson Bailey Haramis Perry Robbins

Underwood Freeman

Runge Sauer Melancon Hatfield

Reconstructed Wetlands of the Tidal Anacostia

| Site | Acres | Date reconstructed |
|------------------------|-------|--------------------|
| 1. Kenilworth Marsh | 31 | 1993 |
| 2. Benning Power Plant | 0.5 | 1996 |
| 3. Kingman Marsh | 40 | 2000 |
| 4. Fringe Marsh | 16 | 2003 |
| 5. Heritage Marsh | 5 | 2006 |
| 6. Bladensburg Marsh | 1 | 2006 |
| 7. Anacostia 11 | 21 | 2007 |



- RECONSTRUCTED ANACOSTIA WETLANDS
2006



Un-dredged

1927

Dredged



- Photo mosaic of Anacostia study wetlands with dates of reconstruction and acreages












08.08.2003 10:47

CONCLUSIONS

1. Vegetation parameters such as cover, richness, diversity, presence of annuals/perennials, etc. were useful for tracking the marsh restoration process and progress. In so doing they could expose such influencing factors as goose herbivory, invasive plant interactions and low sediment elevations.



- Figure 1.** Total vegetative cover. Data points represent least squares means \pm SE. Labels are based on Tukey test results (family-wise error rate $\alpha = 0.05$). Within areas (Fig. 1a), monthly means sharing the same upper-case letters are not significantly different from year to year within the same month. Within a sampling event (Fig. 1b), means sharing the same lower-case letters are not significantly different. Unlabeled series have no significant differences. Reading of the University of Maryland transects at Patuxent Marsh was discontinued after 2004.




Figure 3. Species richness of areas over time.

Data points represent least square means \pm 1 SE. Labels are based on Tukey test results (overall $\alpha = 0.05$). Within areas (Fig. 3a), means sharing the same upper-case letters are not significantly different from year to year. Within a sampling event (Fig. 3b), means sharing the same lower-case letters are not significantly different. Unlabeled series have no significant differences.

2. As measured by laser level there was a significant elevation loss (about 2 inches) at Kingman Area 1 from 2001-2004 and a considerable elevation loss at Kingman Area 2 (1.5") for the same time frame.

| | Transect # | Elevation 2001 feet | Elevation 2004 feet | Sediment Elevation Change Feet | Elevation Change Cm |
|---------------------------|------------|---------------------------|---------------------------|-----------------------------------|------------------------|
| Kingman Area 1 planted | 1 | 1.51 | 1.11 | -0.40 | -12.2 |
| | 2 | 1.37 | 1.15 | -0.22 | - 6.7 |
| | 3 | 1.53 | 1.53 | 0.00 | -0.0 |
| | 4 | 1.93 | 1.73 | -0.20 | -6.1 |
| | 5 | 1.89 | 1.67 | -0.22 | -6.7 |
| | 6 | 2.18 | 2.04 | -0.14 | -4.3 |
| | 7 | 1.86 | 1.68 | -0.18 | -5.5 |
| | 8 | 1.82 | 1.76 | -0.06 | -1.8 |
| | 9 | 1.57 | 1.47 | -0.10 | -3.1 |
| | 10 | 2.02 | 2.05 | +0.03 | +0.9 |
| | 11 | 1.82 | 1.66 | -0.16 | -4.9 |
| unplanted | 1 | 2.25 | 2.14 | -0.11 | -3.4 |
| | 2 | 2.16 | 1.88 | -0.28 | -8.5 |
| | 3 | 1.53 | 1.40 | -0.13 | -4.0 |
| Kingman area 2 planted | 1 | 1.97 | 1.80 | -0.17 | -5.2 |
| | 2 | 1.87 | 1.87 | 0.00 | -0.0 |
| | 3 | 1.64 | 1.48 | -0.16 | -4.9 |

3. Longer periods of inundation at lower elevations reduced the ability of vegetation to rebound from grazing as seedling germination was reduced and plant growth slowed.




Figure 11. Cover by annuals over time.

Data points represent least square means \pm 1SE. Labels are based on Tukey test results (overall $\alpha = 0.05$). Within areas (Fig. 11a), means sharing the same upper-case letters are not significantly different from year to year. Within a sampling event (Fig. 11b), means sharing the same lower-case letters are not significantly different. Unlabeled series had no significant differences.





Figure 7. Cover by annuals at Patuxent over time.

Data points represent least square means \pm 1SE. Labels are based on Tukey test results (overall $\alpha = 0.05$). Within transects, means sharing the same upper-case letters are not significantly different from year to year. Transects lacking labels had no significant year-to-year differences.

4. Data from Surface Elevation Tables (SETs) at Kingman and Kenilworth Marshes documented ongoing vertical accretion in areas not subject to erosion, but no net change in elevation revealed subsidence is still occurring.

Kingman Marsh - Anacostia River

(5 pipe average)



5. Data from hydrologgers revealed higher than normal water levels from greater than normal rainfall in 2003 and 2004 which further hindered revegetation processes by extending periods of inundation.

Water Level Readings for Dueling Creek ('02 vs. '03)




Figure 17. Water levels at Dueling Creek Marsh for the period 5/23-8/7 for 2003 (a wet year) averaged 2-5" higher than for the same period in 2002 (a dry year).

6. Observations from exclosures placed at various elevations revealed the potential for revegetation where grazing was averted and where elevations were suitable to support vegetation growth.





2004.08.13 11:18





2006.08.15

EXCLOSURE STUDY PLOTS



Mod 3/18/04



X SET3

X SET4 1.99



7/17/02



7. There was ample seed dispersal and seed bank available such that conditions of weak vegetative regeneration could not be attributed to these factors.



8. At Kingman Marsh, planted species did not contribute importantly to the reconstructed marsh over time.




Figure 10. Cover by species planted and replanted* at Kingman.

Data points represent least square means \pm 1SE. Labels are based on Tukey test results (overall $\alpha = 0.05$). Within areas, means sharing the same upper-case letters are not significantly different from year to year. Within a sampling event, means sharing the same lower-case letters are not significantly different. Unlabeled series have no significant differences.

Conclusions for Kingman Marsh

- Large functional seed bank
- Vegetation tracked processes
- Emergent vegetation including planted vegetation greatly reduced;
Planted species provide limited cover
- High water levels and sediment loss reduce revegetation
- Sedimentation as well as erosion is occurring; also subsidence. No net elevation change overall
- Marsh impacted from over-abundant geese coupled with lowered elevations;
- Exclosures reveal marsh potential
- Need for some form of resident Canada goose management; EA being written



2006.08.17